



APPENDIX (for online publication only)

Appendix Figure 1 – Differential Changes by Race/Ethnicity in Days of Last 30 Not in Good Mental Health




Notes: Regressions control for race/ethnicity, gender, age, education, number of children, marital and veteran status, and state and quarter-by-year fixed effects. Dots represent point estimates; bars represent 95% confidence intervals. Standard errors are heteroskedasticity-robust and clustered by month-by-year. BRFSS sampling weights are used. Sample sizes are 839,501, 834,801, and 834,730 in Panels A, B, and C.

Appendix Figure 2 – Differential Changes by Race/Ethnicity in Days of Last 30 Drinking Alcoholic Beverages




Notes: Regressions control for race/ethnicity, gender, age, education, number of children, marital and veteran status, and state and quarter-by-year fixed effects. Dots represent point estimates; bars represent 95% confidence intervals. Standard errors are heteroskedasticity-robust and clustered by month-by-year. BRFSS sampling weights are used. Sample sizes are 839,501, 834,801, and 834,730 in Panels A, B, and C.

Appendix Figure 3 – Differential Changes by Race/Ethnicity in Probability of Exercising in the Past 30 Days




Notes: Regressions control for race/ethnicity, gender, age, education, number of children, marital and veteran status, and state and quarter-by-year fixed effects. Dots represent point estimates; bars represent 95% confidence intervals. Standard errors are heteroskedasticity-robust and clustered by month-by-year. BRFSS sampling weights are used. Sample sizes are 839,501, 834,801, and 834,730 in Panels A, B, and C.

Appendix Figure 4 – Differential Changes by Race/Ethnicity in the Probability of Smoking Daily



Notes: Regressions control for race/ethnicity, gender, age, education, number of children, marital and veteran status, and state and quarter-by-year fixed effects. Dots represent point estimates; bars represent 95% confidence intervals. Standard errors are heteroskedasticity-robust and clustered by month-by-year. BRFSS sampling weights are used. Sample sizes are 839,501, 834,801, and 834,730 in Panels A, B, and C.

Appendix Figure 5 – Differential Changes by Race/Ethnicity in the Probability of Using Marijuana in Past 30 Days



Notes: Regressions control for race/ethnicity, gender, age, education, number of children, marital and veteran status, and state and quarter-by-year fixed effects. Dots represent point estimates; bars represent 95% confidence intervals. Standard errors are heteroskedasticity-robust and clustered by month-by-year. BRFSS sampling weights are used. Sample sizes are 839,501, 834,801, and 834,730 in Panels A, B, and C.